

Decision Letter (MJMHS-2021-0007.R1)

From: normala_ib@upm.edu.my

To: santhna@lincoln.edu.my

CC:

Subject: Malaysian Journal of Medicine & Health Sciences - Decision on Manuscript ID MJMHS-2021-0007.R1

Body: 19-Feb-2021

Dear Prof. Dr. Panduragan:

It is a pleasure to accept your manuscript entitled "The Effectiveness Rom (Range Of Motion) Mobilization Education On Informal Caregivers Of stroke Patients In the inpatient Room" in its current form for publication in the Malaysian Journal of Medicine & Health Sciences. The comments of the reviewer(s) who reviewed your manuscript are included at the foot of this letter.

Thank you for your fine contribution. On behalf of the Editors of the Malaysian Journal of Medicine & Health Sciences, we look forward to your continued contributions to the Journal.

Sincerely,

Dr. Normala Ibrahim

Editor-in-Chief, Malaysian Journal of Medicine & Health Sciences

normala_ib@upm.edu.my

Associate Editor Comments to Author:

Associate Editor

Comments to the Author:

(There are no comments.)

Reviewer(s)' Comments to Author:

Reviewer: 1

Comments to the Author

The article is ready for publication.

Reviewer: 2

Comments to the Author

(There are no comments.)



**The Effectiveness Rom (Range Of Motion) Mobilization
Education On Informal Caregivers Of stroke Patients In the
inpatient Room**

Journal:	<i>Malaysian Journal of Medicine & Health Sciences</i>
Manuscript ID	MJMHS-2021-0007.R1
Manuscript Type:	Supp: GHWB
Keywords:	6

SCHOLARONE™
Manuscripts

Table I: Frequency Distribution of Informal Caregiver Gender

Gender	Frequency	Percent age (%)
Male	6	22.2
Female	21	77.8
Amount	27	100.0

Table II: Frequency Distribution of Informal Caregiver Education Level

Education	Frequency	Percentage (%)
SD	7	25.9
SMP	3	11.1
SMA/SMK	10	37.0
D1/D2/D3/D4	3	11.1
S1	4	14.8
Amount	27	100.0

Table III. Distribution of Education and knowledge Levels

Education	Knowledge	
	Effective (≥60)	Ineffective (≤60)
SD	7	
SMP	3	
SMA/SMK	10	
D1/D2/D3/D4	3	
S1	4	
Amount	27	0

Table IV: Wilcoxon Test Analysis

Knowledge	Mean	P Value
Pretest	45.74	0.000
Posttest	80.74	

ABSTRACT

Introduction: Stroke is neurological deficit acute disease attributable to a focal vascular cause known as one of the leading causes of death and disability. Disability caused by stroke also impacts on the family members as their informal caregiver who has to provide the treatment and care with sufficient knowledge. The purpose of this study is to know the effectiveness of the education on ROM (Range of Motion) mobilization to the informal caregivers for stroke patients in the care room of RSUD dr. Soekardjo, Tasikmalaya. **Methods:** The type of this research is quantitative and the analysis was carried out by using non parametric Wilcoxon tTest. The total sample size of this research was 27 respondents, selected using accidental sampling technique. **Result:** The result show a significant differences in the mean value of the research of ROM mobilization education before and after the intervention (p -value < 0.05). **Conclusion:** Therefore, family caregivers should search more information about stroke and its treatment mainly with non-pharmacology component to increase the knowledge in order to minimize stroke recurrence or future complication.

Keywords: Stroke, Informal Caregiver, Education, Range Of Motion

INTRODUCTION

The American Heart Association (AHA, 2014) stated that the incidence of stroke is 3 million stroke patients per year and 50,000 stroke patients with initial stroke occur every year (1). In Indonesia, 500,000 cases were found, with 125,000 of them dying, and the rest had minor or severe disabilities. The death rate due to stroke is estimated to reach 6.5 million patients in 2015 and the number will increase to 8 million patients by 2030 (2). According to 2014 Ministry of Health data, the incidence of stroke is mostly found in the 45-54 year age group. Data for each region in Indonesia, shows West Java 533,895 people (highest), Central Java 431,021 and West Papua 2,007 people (3).

Stroke is one of the leading causes of death and disability worldwide apart from cancer (4). Disability due to stroke can include speech disorders (aphasia) (5) activity disturbances, visual disturbances, (6) weakness, difficulty walking and coordinating body parts, headaches, facial muscle weakness, visual disturbances, sensory disturbances, disturbances in thought processes and loss of control over motor movements (7), which also affect family members (8) as informal caregivers to help the patient's difficulties in completing daily activities, as the patients are very much dependent on emotional and physical support (9).

Informal care is the care given at home and considered non-professional as they do not require any payments from the sufferers. The caregiver's task is to provide emotional support, care for patients (bathe, dress, prepare meals, prepare medicines), manage finances; make decisions about treatment and communicate with formal health services (10).

One of the non-pharmacological treatments that can be given to stroke patients is ROM exercises which can be divided into two, namely active ROM or that done independently, and passive ROM carried out with assistance (11).

A caregiver's attention is very important because the successful treatment and care of stroke patients really needs caregiver's help and support (12), and is expected to improve care for stroke patients to minimize the occurrence of physical disability and dependence of stroke patients in fulfilling their daily needs (13). However, different studies on the role of caregivers in improving the health quality of stroke patients has been carried over but still need more to move forward (14-16).

Therefore, this study aims to identify the effectiveness of ROM mobilization education for informal caregivers for stroke patients in the Inpatient Room of RSUD Dr. Soekardjo, Tasikmalaya, Indonesia.

MATERIALS AND METHODS

This research is a quantitative research, using a Pre-Experimental design in the form of One-Group Pretest-Posttest Design, conducted from 27 April - 25 May 2019. The samples are informal caregiver who accompanies the patient as the person in charge, totaling to 27 respondents. The respondents were selected using Accidental Sampling which was conducted for 4 weeks. The things studied included gender, level of education and the effect of education on mobilization.

The research instrument consisted of 20 numbers related to knowledge about stroke and ROM (Range of Motion) which had to be filled in by respondents who were willing to take part in the research after filling out the informed consent form.

Univariate data analysis performed were through frequency distribution of gender and education Bivariate analysis were carried out using the non-parametric Wilcoxon test which shows the results of the pretest-posttest value with a significance value *p* value of <0.05.

RESULTS

The results of this research regarding the characteristics of respondents were obtained from the results of research on the Effectiveness of ROM (Range of Motion) Mobilization Education on Informal Caregivers of Stroke Patients in the Inpatient Room of RSUD dr. Soekardjo, Tasikmalaya are as follows:

1) Gender

Table I: Frequency and percentage Distribution of Informal Caregiver Gender

Based on table I the female caregivers were more common with a total of 21 people (77.8%) and male caregivers with a total of 6 people (22.2%)

2) Education

Table II: Frequency and Percentage Distribution of the Education Level of Informal Caregiver

Based on table II the level of education of the respondents show SD (Sekolah Dasar) education as many as 7 respondents (25.9%), SMP(Sekolah Menengah Pertama) 3 (11.1%), and the dominant education level of the respondents in this study was the SMA(Sekolah Menengah Atas) / SMK(Sekolah Menengah Kejuruan) / equivalent level as many as 10 people (37.0%), D1(Diploma1), D2(Diploma2), D3(Diploma3), D4(Diploma4) as many as 3 respondents or (11.1%) and S1(Sarjana S1) as many as 4 respondents or (14.8%).

3). Education&Knowledge

Table III. Distribution of Education and knowledge Levels

TableIII shows that the respondents in this study showed a value above 60 or effective, which means that education or counseling at all levels of education, is at an effective value.

4). Wilcoxon Test Analysis

Table IV: Wilcoxon Test Analysis

Table IV shows the value of the results of the non-parametric test analysis Wilcoxon Test. Based on the statistical output the average pretest value is 45.74 which shows the average is low, while the posttest average value is 80.74 which shows the average value is high. The educational effectiveness value of ROM Mobilization on Informal Caregiver of Stroke Patients in the Inpatient Room of RSUD Dr. Soekardjo,Tasikmalaya shows that the p - value is 0.000, (p -value < 0.05) and thus H_a is accepted and there is a significant difference in the average of before and after the ROM mobilization education.

DISCUSSION

The results of this study indicate that female respondents are more than male respondents with the highest level of education is SMA / SMK / SLTA / equivalent, and the level of knowledge is in the low range.

Respondents in the study who are informal caregivers have a duty to support other individuals in emotional support, caring for patients, managing finances, making decisions about treatment and communicating with formal health services (10), really need good knowledge.

Knowledge of ROM (Range of Motion) which is a therapy to maintain or improve the level of perfection of the ability to move joints normally and completely to increase muscle mass and

muscle tone, which is included in non-pharmacological therapy is still common to respondents or their families (17).

Respondents' initial knowledge can be categorized as low but there is a significant difference after the respondent received education from the researcher as evidenced by the pretest and posttest scores.

The knowledge that is needed by each individual can not only be obtained through formal education as-, there are many other ways to get knowledge which include through counseling to exchange knowledge, social media can also be a means of obtaining information related to stroke or other information and can be obtained through sight and hearing.

The results of this study are in line with the research (18) which stated that family knowledge about stroke is mostly in the sufficient category, namely 49% and there is a relationship between knowledge and family attitudes in providing care for post-stroke patients at the Neurology Polyclinic RSUD dr. Soekardjo, Tasikmalaya.

Another research suggests that knowledge can be obtained through the five senses, especially sight and hearing. Sensing occurs through the five human senses, namely the senses of sight, hearing, smell, taste, sense of touch, and most significantly by the eyes and ears, which are the most easiest media to gain knowledge (19).

The results of the measurement of knowledge showed that there was no decrease in the posttest score, instead there was an increase after education was given which could support informal caregivers to provide care to sick family members.

Another research which states that the caregiver's experience in caring for post-stroke patients have a positive impact. The positive impact is in the form of increasing self-understanding of the meaning of life, increasing knowledge and experience and repaying favors as a form of obligation as a family (20).

This is evidenced by the posttest scores and short interviews with respondents after being given education about ROM mobilization which states that they are more familiar with strokes and are now aware of the method of treatment for family members when they come home from the hospital later to prevent complications.

Many stroke patients experience recurrent strokes or complications after discharge from hospital which actually worsens the patient's condition because the care by the informal

caregiver or family while at home is not good because of poor knowledge about the patient's illness and how to treat it.

This is reinforced by the results of research which states that the level of respondent's knowledge with the respondent's attitude in providing care to post-stroke patients has a significant relationship (21).

The results showed the effectiveness of providing education to caregivers as seen from the p value of 0.000 which is smaller than the value of α 0.05 which means there is a significant difference before and after being given education in the inpatient room of RSUD Dr. Soekardjo, Tasikmalaya.

After being given education about passive or active ROM, the informal caregiver and the patient are excited to do ROM at home after being discharged from the hospital. The caregiver feels helped by the education given because it can increase knowledge and make it easier for the caregiver to care for family members who have incurred stroke. Stroke patients also feel better after provided with family-assisted ROM mobilization during the counseling session because this helped to stretch the affected muscles.

This is in line with the research conducted by Semet *et al.* (22) that proved that patient families had a high level of knowledge about stroke and the relatives of patients had high level of knowledge of the stroke. Knowledge of stroke patients was higher than their relatives

CONCLUSION

Analysis of the results of the pretest level of knowledge of the informal caregiver of stroke patients showed low results, which showed that the respondents' level of initial knowledge was in the low category ≤ 60 , while the results of the post test showed a significant increase in the level of knowledge of the respondents after being given education in the high category ≥ 60 . Wilcoxon non-parametric test analysis of the effectiveness of ROM (Range of Motion) mobilization education on informal caregivers of stroke patients showed effective results with a p value of 0.000 ($0.000 < 0.05$).

SUGGESTION

For informal caregivers, hopefully it can be an input in increasing knowledge about stroke by attending stroke-related counseling or by reading different sources of information to obtain

additional knowledge for education. The information obtained from this study can become a source for further research development.

ACKNOWLEDGMENT

All authors are very thankful to the management committee of Doctor Soekardjo Regional General Hospital for allowing authors to collect data. The authors are also thankful to Stikes Muhammadiyah Ciamis for the continuous support to complete this research work.

For Review Only

REFERENCE

1. Goldstein L. Guideline for the Primary Prevention of Stroke. A Guideline for Health care Professionals from the American Heart Association. American Stroke Association. 2014.
2. Kusuma Y, Kiemas L, Misbach J. Burden of stroke in Indonesia. *Int J Stroke*. 2009;4(5):379-80.
3. Kemenkes RI. Infodatin : Situasi Kesehatan Jantung. Pus Data dan Inf Kementerian Kesehatan RI [Internet]. 2014;1–8.
4. Bartoli F, Lillia N, Lax A, Crocamo C, Mantero V, Carrà G, Agostoni E, Clerici M. Depression after stroke and risk of mortality: a systematic review and meta-analysis. *Stroke research and treatment*. 2013 Oct;2013.
5. Haryanto GD, Setyawan D, Kusuma MA. Pengaruh Terapi AIUEO terhadap kemampuan bicara pada pasien stroke yang mengalami afasia motorik di RSUD Tugurejo Semarang. *Karya Ilmiah*. 2014 Jun 15.
6. Lee A. Understanding patients and their spouse's experiences of home rehabilitation after a stroke. *International Journal of Integrated Care*. 2011 Jul;11(Suppl).
7. Sari SH, Agianto A, Wahid A. Batasan Karakteristik Dan Faktor Yang Berhubungan (Etiologi) Diagnosa Keperawatan: Hambatan Mobilitas Fisik Pada Pasien Stroke. *Dunia Keperawatan: Jurnal Keperawatan dan Kesehatan*. 2015 Mar 1;3(1):12-21.
8. A'la MZ, Yosep I, Agustina HR. Pengaruh bereavement life review terhadap kesejahteraan spiritual pada keluarga pasien stroke. *Jurnal Keperawatan Padjadjaran*. 2017;5(2).
9. Fadilla L. Peran Harapan Dan Resiliensi Terhadap Stres, Kecemasan Dan Depresi Pada Caregiver Penderita Stroke (Doctoral dissertation, Universitas Islam Negeri Sultan Sarif Kasim Riau).
10. Putri AK. EKSPRESI EMOSI PENDAMPING SKIZOFRENIA (Doctoral dissertation, UIN Sunan Ampel).2015
11. Wijanarko MO, Setyawan D, Kusuma MA. Pengaruh Terapi Musik Klasik Terhadap Pasien Stroke Yang Menjalani Latihan Range of Motion (ROM) Pasif. *Karya Ilmiah*. 2014 Jun 15.
12. Daulay NM, Setiawan S, Febriany N. Pengalaman Keluarga sebagai Caregiver dalam Merawat Pasien Strok di Rumah. *Jurnal Keperawatan Padjadjaran*. 2014;2(3).
13. Festy P. Peran keluarga dalam pelaksanaan rehabilitasi medik pada pasien stroke. *Jurnal Penelitian Fakultas Ilmu Kesehatan Universitas Muhammadiyah Surabaya*. 2012.
14. Choong K, Canci F, Clark H, Hopkins RO, Kudchadkar SR, Lati J, Morrow B, Neu C, Wiecezorek B, Zebuhr C. Practice Recommendations for Early Mobilization in Critically Ill Children. *J Pediatr Intensive Care*. 2018 Mar;7(1):14-26. doi: 10.1055/s-0037-1601424. Epub 2017 Apr 10. PMID: 31073462; PMCID: PMC6260323.

1
2
3 15. King RB, Ainsworth CR, Ronen M, Hartke RJ. Stroke caregivers: pressing problems reported
4 during the first months of caregiving. *J Neurosci Nurs*. 2010 Dec;42(6):302-11. doi:
5 10.1097/jnn.0b013e3181f8a575. PMID: 21207768; PMCID: PMC3064495.
6
7 16. Baatiema L, de-Graft Aikins A, Sarfo FS, Abimbola S, Ganle JK, Somerset S. Improving the
8 quality of care for people who had a stroke in a low-/middle-income country: A qualitative
9 analysis of health-care professionals' perspectives. *Health Expect*. 2020 Apr;23(2):450-460. doi:
10 10.1111/hex.13027. Epub 2020 Jan 22. PMID: 31967387; PMCID: PMC7104640.
11
12 17. Agusrianto A, Rantesigi N. Penerapan Latihan Range of Motion (ROM) Pasif terhadap
13 Peningkatan Kekuatan Otot Ekstremitas pada Pasien dengan Kasus Stroke. *Jurnal Ilmiah*
14 *Kesehatan*. 2020;2(2):61-6.
15
16 18. Kustiawan R, Badruzaman R. Hubungan Tingkat Pengetahuan Dengan Sikap Keluarga Dalam
17 Pemberian Perawatan Pasien Pasca Stroke Di Poliklinik Syaraf Rsud Dr. Soekardjo Kota
18 Tasikmalaya. *Jurnal Kesehatan Bakti Tunas Husada: Jurnal Ilmu-ilmu Keperawatan, Analisis*
19 *Kesehatan dan Farmasi*. 2016 Nov 11;15(1):92-9.
20
21 19. Notoatmodjo S. Promosi kesehatan teori dan aplikasi. Jakarta: Rineka Cipta. 2005:52-4.
22
23 20. Opara JA, Jaracz K. Quality of life of post-stroke patients and their caregivers. *Journal of*
24 *medicine and life*. 2010 Aug 15;3(3):216.
25
26 21. Sonatha B, Gayatri D. Hubungan tingkat pengetahuan dengan sikap keluarga dalam pemberian
27 perawatan pasien pasca stroke. Universitas Indonesia, Depok. 2012 Jun.
28
29 22. Semet GR, Kembuan MA, Karema W. Gambaran pengetahuan stroke pada penderita dan
30 keluarga di RSUP Prof. Dr. RD Kandou Manado. *e-CliniC*. 2016;4(2).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60